

Technical Bulletin

Genes that fit *your* farm.

SeCan

Canada's Seed Partner

CDC Hickie Yellow Field Pea



Progress Through Research
Le progrès grâce à la recherche

Description:

CDC Hickie is a high yielding yellow cotyledon field pea with high protein, medium vine length, good standability and good resistance to seed coat breakage, dimpling and less green coloration.

Parentage: CDC 2283-17/P0410-16

Strengths:

- 104% yield of CDC Amarillo
- +0.2% protein compared to CDC Amarillo
- Good lodging resistance; similar to CDC Amarillo and AAC Lacombe
- Medium-long vine length; similar to CDC Amarillo
- Powdery mildew resistance
- Moderate resistance rating to Fusarium wilt
- Improved seed characteristic package- rated good for:
 - Resistance to seed coat breakage
 - Resistance to dimpling

- Greenness (less green coloration) better than CDC Inca or CDC Spectrum

Neutral Traits:

- Medium maturity, 1 day later than CDC Amarillo
- Medium seed size similar to CDC Amarillo
- Fair Mycosphaerella blight resistance (similar to CDC Amarillo or CDC Lewochko)

PBR Protected

Breeder:

Funded by:



UNIVERSITY OF SASKATCHEWAN
Crop Development Centre
COLLEGE OF AGRICULTURE AND BIORESOURCES
AGBIO.USASK.CA



Table 1: Performance of CDC Hickie compared to the checks (Pea Co-op Test B, 2017-2018)

Variety	Yield (kg/ha)	Yield (% mean of checks)	Days to maturity	Vine length (cm)	Lodging (pre-harvest)	Seed wt (mg)	Shape	Crude protein (%)
CDC Amarillo (check)	3900	100	93	83	2.6	229	2.5	24.2
AAC Lacombe (check)	3930	100	92	80	2.5	256	2.5	22.7
CDC Hickie	4040	103	94	83	2.6	232	2.7	24.4
# of locations	27		25	21	25	26	18	16

Lodging score: 1=upright; 9=completely lodged
Shape: 1=round; 5=cubic

Table 2: Performance of CDC Hickie compared to the checks (Pea Co-op Test B, 2017-2018)

Variety	Myc (AB)	Myc (SK)	Fusarium wilt (%)	Powdery mildew
CDC Amarillo (check)	3.2	4.7	29	0
AAC Lacombe (check)	2.1	4.9	19	0
CDC Hickie	1.8	5.1	18	0
# locations	2	2	2	2

Myc (MB)= Mycosphaerella blight assessed at Morden, MB (1=no disease, 9=completely blighted)

Myc (SK)= Mycosphaerella blight assessed at Saskatoon, SK.

Powdery mildew (0=no disease, 9=completely mildewed)

For more information, call 1-800-665-7333 or visit www.secan.com

2025 Seed Manitoba

Variety	Yield % of Check AAC Carver	Site Years Tested	Relative Maturity	Relative Vine Length	Seed Size (TKW)	Resistance to:						
						Lodging	Powdery Mildew	MB ¹	Fusarium Wilt ²	Green Seed Coat ³	Seed Coat Breakage	Seed Coat Dimpling ⁴
AAC Carver	100	50	Early	L	240	G	VG	F	I	G	G	G
CDC Boundless	97	9	Mid	L	230	VG	VG	F	MR	G	G	G
CDC Lewochko	100	42	Long	L	230	VG	VG	F	I	G	G	G
CDC Tollefson	104	23	Long	L	240	VG	VG	F	MR	G	G	G
CDC Hickie	98	23	Mid	M	230	VG	VG	F	MR	G	G	G

1. Mycosphaerella Blight
2. Varieties which show good disease tolerance to one strain of Fusarium wilt may be susceptible to other strains
3. Green seed coats: G= 0-10%, F= 11-25%
4. Seed coat dimpling rating: VG= 0-5%, G= 6-20%, F= 21-50%

2025 Saskatchewan Seed Guide

Variety	Years Tested	Yield (% of CDC Amarillo)			Protein	Relative Maturity	Lodging ¹	Vine Length (cm)	Resistance to:						Seed Weight (g/1000)
		1, 2, & South 3	North 3 & 4	Irrigation					MB ²	Powdery Mildew	Fusarium Wilt	Seed Coat Breakage	Seed Coat Dimpling ³	Greenness ⁴	
CDC Amarillo (C)	15	100	100	100	23.8	M	2.9	85	3.9	R	MR	F	F	G	230
CDC Boundless	5	109	105	---	+0.5	M	-0.1	90	-0.1	R	MR	G	G	G	230
CDC Canary	10	99	100	---	-0.2	E	+0.2	85	+0.4	R	I	G	F	F	230
CDC Lewochko	11	103	103	---	+0.7	M	-0.1	90	-0.1	R	I	G	G	G	230
CDC Spectrum	13	104	103	93	+0.3	M	0.0	85	-0.2	R	I	G	G	F	240
CDC Tollefson	8	107	106	---	-0.1	M	-0.2	90	-0.1	R	MR	G	G	G	240
CDC Hickie	8	106	105	---	+0.6	M	-0.1	85	0.0	R	MR	G	G	G	230

1. Lodging: 1= completely upright, 9= completely lodged; values are +/- CDC Amarillo
2. Mycosphaerella Blight score (1-9): 1= no disease, 9= completely blighted, values are +/- CDC Amarillo
3. Seed coat dimpling: VG= 0-5%, G= 6-20%, F= 21-50%
4. Green seed coats: G= 0-15%, F= 16-40%

2025 Alberta Seed Guide

Variety	Most Recent year of RVT Testing	Overall Yield	Overall Station Years of Testing	Brown - Irrigated		Brown		Black-short		Black-mid		Grey Wooded		Agronomic Characteristics:					Disease Tolerance ⁵		Tolerance to:			
				Yield (%)	Site Years	Yield (%)	Site Years	Yield (%)	Site Years	Yield (%)	Site Years	Yield (%)	Site Years	Yield (%)	Site Years	Protein (%)	Maturity Rating ¹	Vine Length (cm)	Seed Size (TKW)	Standability ²	MB ³	Fusarium Wilt	Seed Coat Breakage	Seed Coat Dimpling ⁴
CDC Amarillo (bu/ac)		74		92		64		80		69		76												
CDC Amarillo (C)	2024	100	87	100	4	100	10	100	22	100	38	100	13	23.8	M	85	230	2.9	3.9	MR	F	F	G	
CDC Canary	2023	103	58	XX	3	95	7	104	14	105	23	103	13	-0.2	E	58	230	3.1	4.3	I	G	F	F	
CDC Lewochko	2023	103	58	XX	3	98	7	104	14	104	23	103	11	+0.7	M	90	230	2.8	3.8	I	G	G	G	
CDC Spectrum	2023	103	58	XX	3	94	7	105	14	105	27	103	11	+0.3	M	85	240	2.9	3.7	I	G	G	F	
CDC Tollefson	2024	105	25	XX	2	XX	2	108	6	106	10	XX	5	-0.1	M	90	240	2.7	3.8	MR	G	G	G	
CDC Hickie	2024	101	25	XX	2	XX	2	105	6	101	10	XX	3	+0.6	M	85	230	2.8	2.9	MR	G	G	G	

1. Maturity: E= Early, M= Medium, L= Late
2. Standability: 1= completely upright, 9= completely lodging
3. Mycosphaerella Blight: 1= no disease, 9= completely blighted
4. Seed Coat Dimpling: VG= 0-5%, G= 6-20%, F= 21-50%
5. Disease Tolerance: I= intermediate, MR= moderately resistance, R= resistant
6. Green Seed Coat: G= 0-10%, F= 11-25%

For more information, call 1-800-665-7333 or visit www.secan.com